

FISH SURVEY REPORT

Lake Santee

December 3, 2018

**Prepared for:
Lake Santee P.O.A. Inc.
13 South Wrenn Parkway
Greensburg, IN, 47240**

**Prepared by:
Aquatic Control Inc.
P.O. Box 100
Seymour, IN 47274**

Table of Contents

Introduction	1
Results and Discussion	1
Water Chemistry	1
Fish Collection	2
Bluegill.....	3
Gizzard Shad	6
Black Crappie.....	8
Largemouth Bass.....	9
Brown Bullhead	12
Channel Catfish.....	12
Other species	13
Summary and Recommendations.....	15
Appendix A.....	17
General Information	17
Equipment and Methods	20
Literature Cited and Reference List	22
Appendix B-Fish Collection Table	25

List of Figures

Figure 1. Temperature and dissolved oxygen profiles for Lake Santee	2
Figure 2. Relative abundance of species collected.....	3
Figure 3. Photograph of bluegill	4
Figure 4. Bluegill length frequency.....	4
Figure 5. Bluegill relative abundance over the past three surveys	5
Figure 6. Bluegill standard weight comparison	5
Figure 7. Photograph of Gizzard Shad.....	6
Figure 8. Gizzard Shad length frequency.....	7
Figure 9. Gizzard Shad relative abundance (4.0-9.5 in.) past three surveys	7
Figure 10. Gizzard Shad relative abundance (10.0-15.5 in.) past three surveys	8
Figure 11. Gizzard Shad CPUE past four surveys	8
Figure 12. Photograph of Black Crappie.....	9
Figure 13. Photograph of Largemouth Bass.....	9
Figure 14. Largemouth Bass length frequency	10
Figure 15. Largemouth Bass relative abundance past three surveys.....	10
Figure 16. Largemouth Bass relative abundance past three surveys.....	11
Figure 17. Largemouth Bass standard weight comparison	11
Figure 18. Photograph of Brown Bullhead	12
Figure 19. Photograph of Channel Catfish	13
Figure 20. Photograph of Redear Sunfish.....	13
Figure 21. Photographic comparison of Bluegill and Green Sunfish	14
Figure 22. Photograph of Hybrid Striped Bass	14

List of Tables

Table 1. Water quality data.....	2
Table 2. Species collected.....	3

INTRODUCTION

A Standard Survey of the fish community and other physical, biological, and chemical factors directly affecting the fish community was completed at Lake Santee on October 8, 2018. The major objectives of this survey and report are:

1. To provide a current status report on the fish community of the lake.
2. To compare the current characteristics of the fish community with established indices and averages for Indiana lakes and with past surveys on Lake Santee.
3. To provide recommendations for management strategies to enhance or sustain the sport fish community.

The data collected are adequate for the intended uses; however, there will be unanswered questions regarding aspects of the fish population and other related factors of the biological community in the lake. All fish numbers used in the report are based on the samples collected and should not be interpreted to be absolute or estimated numbers of fish in the lake. General information regarding water chemistry, fish communities, and methods are described in Appendix A. A detailed fish collection table is presented in Appendix B.

RESULTS AND DISCUSSION

WATER CHEMISTRY

The results of selected physio-chemical parameters from Lake Santee are presented in Table 1. Water temperatures ranged from 75.6 degrees Fahrenheit at the surface to 67.4 degrees Fahrenheit at the bottom. Dissolved oxygen ranged from 14.01 parts per-million (ppm) at the surface to 0.22 ppm at the bottom (Figure 1). A desirable oxygen level for maintenance of healthy stress free fish was present to a depth of 12.0 feet. These numbers indicate Lake Santee was stratified at the time of the survey, which is typical for this time of year (See Appendix A for further details on lake stratification). The alkalinity level was 188.1 ppm at the surface and 171.0 at the bottom. The hardness level was 136.8 ppm at the surface and 171.0 at the bottom. The pH was 8.8 at the surface and 7.4 on the bottom. These numbers are normal for lakes in this area and indicate the lake is capable of good fish production. The Secchi disk depth was measured at 2.5 feet. Total nitrogen levels were 1.27 ppm at the surface and 3.45 on the bottom. Total phosphorus levels were 0.17 ppm at the surface and 0.41 ppm on the bottom. Lake Santee appears to have water quality which is capable of supporting a healthy fish population.

Table 1. Selected water quality parameters measured on Lake Santee, October 8, 2018.

Sample Depth (ft.)	Temp.(°F)	Dissolved Oxygen (ppm)	pH (standard units)	Total Alkalinity (ppm)	Total Hardness (ppm)	Total Nitrogen (ppm)	Total Phosphorus (ppm)
Surface	75.6	14.01	8.8	188.1	136.8	1.27	0.17
3	75.5	13.88	-	-	-	-	-
6	74.0	8.42	-	-	-	-	-
9	73.2	6.80	-	-	-	-	-
12	72.5	5.01	-	-	-	-	-
15	71.6	1.91	-	-	-	-	-
18	70.3	0.36	-	-	-	-	-
21	68.2	0.26	-	-	-	-	-
22	67.4	0.22	7.4	171.0	136.8	3.45	0.41




Figure 1. Temperature and dissolved oxygen profiles for Lake Santee, October 8, 2018.

FISH COLLECTION

A total of 1192 fish weighing 248.56 pounds and representing ten species was collected from Lake Santee (Table 2 & Figure 2). Bluegill *Lepomis macrochirus* was the most abundant species comprising 66.53% of the fish collected. Gizzard Shad *Dorosoma cepedianum* was the second most abundant species (12.33%), followed by Black Crappie *Pomoxis nigromaculatus* (9.65%), Largemouth Bass *Micropterus salmoides* (4.87%), Brown Bullhead *Ameiurus nebulosus* (3.27%), Channel Catfish *Ictalurus punctatus* (2.85%), Redear Sunfish *Lepomis microlophus* (0.17%), Green Sunfish *Lepomis cyanellus* (0.17%), Hybrid Striped Bass *Morone chrysops* X *Morone saxatilis* (0.08%),

and Spotfin Shiner *Cyprinella spiloptera* (0.08%). All of these species are desirable in a lake of this size with the exception of Gizzard Shad, Brown Bullhead, and Green Sunfish.

Table 2. Species collected from Lake Santee, October 8, 2018.

Species	N	%N	Size Range (in)	Total Weight (lbs)	%Wt.	PSD	N/hr.
Bluegill	793	66.53%	<3.0-7.5	86.04	34.62%	39	397
Gizzard Shad	147	12.33%	5.5-15.0	30.76	12.38%		74
Black Crappie	115	9.65%	4.0-8.5	17.59	7.08%		58
Largemouth Bass	58	4.87%	3.5-20.0	70.90	28.52%	63	29
Brown Bullhead	39	3.27%	5.0-10.5	14.18	5.70%		20
Channel Catfish	34	2.85%	3.0-20.0	18.36	7.39%		17
Redear Sunfish	2	0.17%	3.5-8.0	0.50	0.20%		1
Green Sunfish	2	0.17%	3.5-5.5	0.13	0.05%		1
Hybrid Striped Bass	1	0.08%	27.0	10.09	4.06%		1
Spotfin Shiner	1	0.08%	<3.0	0.01	0.00%		1
Total	1192			248.56			




Figure 2. Relative abundance of species collected from Lake Santee, October 8, 2018.

Bluegill

Bluegill (Figure 3) was the most abundant species collected (66.53%) and ranked second by weight (24.77%). Individuals ranged in size from less than 3.0 to 7.5 inches (Figure 4). Approximately 5% of Bluegill collected were 3.5 inches or less, indicating poor

reproduction occurred in 2018. There was a large number of quality Bluegill collected compared to young of year. This led to a proportional stock density of 39, which is at the upper end of the desired range of 20-40 for Bluegill (proportion of quality fish within a population, see Appendix A). The PSD for Bluegill has decreased dramatically from 76 in the 2015 survey. The CPUE (Catch-Per-Unit-Effort) was 397 fish/hour compared to 279 fish/hour in the previous survey. The relative abundance of Bluegill collected over the past three surveys is displayed in Figure 5. Condition factors (measurement of overall plumpness) were below average for most size ranges. Bluegill weights were also found to be below standard weights in most size ranges (Figure 6).



Figure 3. Photograph of Bluegill, *Lepomis macrochirus*.




Figure 4. Length frequency distribution of Bluegill collected from Lake Santee, October 8, 2018.




Figure 5. Relative abundance of Bluegill collected from Lake Santee over the past three surveys.




Figure 6. Comparison of Lake Santee Bluegill weights to standard Bluegill weights.

Gizzard Shad

Gizzard Shad (Figure 7) was the second most abundant species collected (12.33%), and ranked third by weight (12.38%). Individuals ranged in size from 5.5 to 15.5 inches (Figure 8). Figures 9 and 10 illustrate the relative abundance of Gizzard Shad at 0.5 length groups over the past three surveys. The CPUE for this survey was 74, compared to 102 (2015), 163 (2012), 138 (2007), and 102 (2005) (Figure 11). As observed in previous surveys, Bluegill don't grow well in lakes containing this species. This may be due to competition with Gizzard Shad for food and space or Largemouth Bass switching to Gizzard Shad as their primary forage causing Bluegill to become overabundant and slow growing. Largemouth Bass typically grow well in the lakes containing Gizzard Shad, especially individuals that are 15.0 inches and larger; however, Gizzard Shad grow rapidly and reach sizes that bass cannot eat. Small shad also negatively affect desirable species through competition for food resources and habitat. This species has a very high reproductive potential. The Gizzard Shad population should continue to be monitored closely, and predator stockings should also continue in an attempt to prevent the shad population from growing larger.



Figure 7. Photograph of Gizzard Shad, *Dorosoma cepedianum*.




Figure 8. Length frequency distribution of Gizzard Shad collected from Lake Santee, October 8, 2018.




Figure 9. Relative abundance of Gizzard Shad (4.0-9.5 in.) collected from Lake Santee over the past three surveys.




Figure 10. Relative abundance of Gizzard Shad (10.0-15.5 in.) collected from Lake Santee over the past three surveys.




Figure 11. Gizzard Shad CPUE over the past five surveys.

Black Crappie

Black Crappie (Figure 12) was the third most abundant species collected (9.65%) and ranked fifth by weight (7.08%). They ranged in size from 4.0 to 8.5 inches.

Approximately 67% of Black Crappie collected were 7.0 to 8.0 inches. The CPUE for this species was 58 fish/hour compared to 41 fish/hour in 2015. It appears that the Black Crappie population may have increased since the last survey; however, crappie inhabit

deeper water and are usually not well represented in electrofishing surveys, so the crappie population is most likely larger than indicated by the survey.



Figure 12. Photograph of Black Crappie, *Pomoxis nigromaculatus*.

Largemouth Bass

Largemouth Bass (Figure 13) was the fourth most abundant species collected (4.87%) and ranked second by weight (28.52%). A total of 58 Largemouth Bass ranging in size from 3.5 to 20.0 inches was collected (Figure 14). A small number of Largemouth Bass less than 7.5 inches were collected indicating reproduction/recruitment continues to suffer but is not absent completely. The sample of Largemouth Bass collected was distributed well across most size ranges. This led to a PSD of 63 for Largemouth Bass, which is on the upper end of the desired range of 40-60. Figures 15 and 16 shows the relative abundance by size class from the past three surveys. The Largemouth Bass collected appeared to be in very good shape. Condition factors (measurement of overall plumpness) were excellent for most size classes. The average weights for the Largemouth Bass were also good compared to standard weights (Figure 17).



Figure 13. Photograph of Largemouth Bass, *Micropterus salmoides*.




Figure 14. Length frequency distribution of Largemouth Bass collected from Lake Santee, October 8, 2018.




Figure 15. Largemouth Bass relative abundance comparison over the past three surveys.




Figure 16. Largemouth Bass relative abundance comparison over the past three surveys.




Figure 17. Comparison of Lake Santee Largemouth Bass weights to standard Largemouth Bass weights.

Brown Bullhead

Thirty-nine Brown Bullheads (Figure 18) were collected while sampling. Brown Bullheads are considered undesirable and compete with desirable species for food resources. This species isn't usually considered an important game fish. They don't reach sizes comparable to Channel Catfish, and most anglers don't find them very palatable. All bullheads that are caught should be removed from the lake.



Figure 18. Photograph of Brown Bullhead, *Ameiurus nebulosus*.

Channel Catfish

Thirty-four Channel Catfish (Figure 19) were collected. Individuals ranged from 3.0 to 20.0 inches. This species is not typically sampled well with electrofishing equipment; however, channel catfish were well represented during the current survey. Young-of-the-year channel catfish are typically eliminated by Largemouth Bass and other predators leading to little or no recruitment into the population. Channel catfish should continue to be stocked periodically if they are desired in the fishery. This species is omnivorous and opportunistic in feeding. They consume many different types of living and dead material. Channel Catfish will aide in putting pressure on overabundant populations of prey species.



Figure 19. Photograph of Channel Catfish, *Ictalurus punctatus*.

Other species

Four other species were collected in low numbers; two Redear Sunfish, two Green Sunfish, one Hybrid Striped Bass, and one Spotfin Shiner. Redear Sunfish (Figure 20) are still present, but continue to be in low numbers. Green Sunfish look superficially like bluegill. They can easily be distinguished by their larger mouths and more rounded pectoral fins. Figure 21 illustrates the differences in appearance between bluegill and green sunfish. Hybrid Striped Bass (Figure 22) were collected in low numbers but have but are likely much more abundant than shown in the survey. Stockings should continue to keep Gizzard Shad populations from growing.



Figure 20. Photograph of Redear Sunfish, *Lepomis microlophus*.



Figure 21. Photographic comparison of Bluegill (top) and Green Sunfish (bottom), *Lepomis cyanellus*.



Figure 22. Photograph of Hybrid Striped Bass, *Morone chrysops* X *Morone saxatilis*.

SUMMARY AND RECOMMENDATIONS

Lake Santee should continue to provide good angling opportunities for trophy Largemouth Bass, Bluegill, Black Crappie, Channel Catfish, and an occasional Hybrid Striped Bass. The Bluegill population continues to be dominated by adult individuals. Predator stockings should result in further reduction of the gizzard shad population and the Bluegill population should continue to improve with lowered Gizzard Shad numbers. Largemouth Bass collected appear very healthy and are likely experiencing rapid growth rates; however, reproduction and recruitment continues to suffer. The supplemental stockings that have taken place each year since the previous survey appear to be sustaining the population. If these stockings do not continue, the Largemouth Bass population may not be able to sustain itself. Largemouth Bass supplemental stockings should continue to be the highest management priority. Structure is still lacking throughout the lake. If individual lot owners were to add structure around their personal docks it could greatly benefit the fishery as a whole by allowing small Largemouth Bass a place to avoid predation after they hatch. Adding structure near a dock will also attract fish to the area that anglers could take advantage of.

The following recommendations, **listed in order of importance**, will help protect and enhance the fishery in Lake Santee:

1. Stock 4,000 5.0 to 8.0 inch Largemouth Bass per year for the next two years. Supplemental stockings are needed to sustain and increase the population due to very poor natural recruitment.
2. Encourage “catch and release” only on Largemouth Bass. This can be adjusted for tournaments, but tournament fishermen should exercise extreme caution when handling fish, especially during hot summer months.
3. Continue stocking Hybrid Striped Bass at a rate of 2,000 per year. Restrictions should continue with a 1 fish per day, 21.0 inch minimum length limit. This will continue to ensure that there is a healthy population in the lake and aid in decreasing Gizzard Shad.
4. No restrictions are necessary on Bluegill harvest. Harvest of this species is encouraged.
5. Continue with the addition of artificial structure. Poor light penetration continues to limit the amount of submerged vegetation. Artificial structure should be the means of replacing this vegetation. If individual lot owners

are interested in artificial structure they can call Aquatic Control for prices.

6. Conduct a Standard Fish Survey in 2021 in order to monitor the effects of the above recommendations and assess needs for further management activities.
7. Continue to work with local landowners in the watershed and encourage the utilization of best land management practices concerning farming techniques. Examples of these practices are the creation of permanent wetlands, no-till farming, or the installation of filter strips.
8. Continue with nuisance bluegreen algae bloom control with targeted EPA algaecide applications. Applications to control problematic bluegreen algae blooms aid in increased water clarity as well as the reduction of extreme dissolved oxygen fluctuations that have been a persistent problem at Lake Santee in the past. Nuisance bluegreen algae blooms also affect juvenile fish. Juvenile fish feed on zooplankton. Bluegreen algae is not a preferred food item for these very beneficial zooplankton. Reduction in Gizzard Shad numbers could also help reduce the severity of bluegreen algae blooms.
9. Maintain the limit Redear Sunfish harvest to 15 per day.
10. Continue the limit on Black Crappie harvest to 25 per day
11. Continue to remove all Brown Bullheads, Green Sunfish, and Common Carp when caught.
12. If Channel Catfish are desired, continue to stock every 2-3 years.

Prepared by: Aquatic Control Inc.
Wesley Goldsmith, Fisheries Biologist

APPENDIX A

GENERAL INFORMATION

In order to help understand our analysis and recommendations, basic principles of water chemistry and the physical attributes of water must be understood. Sources of dissolved oxygen (D.O.) in water include uptake from the atmosphere and photosynthesis.

Decreases in D.O. are mostly attributed to the respiration of plants, animals, and aerobic bacteria that occur in a lake or pond. Large quantities of plants may produce oxygen depletion during the nighttime hours as plants stop photosynthesis and utilize oxygen for respiration. Dissolved oxygen levels below 5.0 are considered undesirable in ponds and lakes (Boyd, 1991). Lower levels of D.O. may stress fish and decrease the rate of decomposition of organic matter entering or produced within a lake or pond. If oxygen depletion is determined to be a problem in a lake or pond, solutions exist to help improve conditions. Vegetation control to reduce overly abundant vegetation may improve conditions. Aeration systems may also be used to increase oxygen levels and promote the breakdown of organic matter.

Water temperature of a lake or pond affects the activity of "cold-blooded" animals such as fish and invertebrates as well as the amount of D.O. that water is capable of holding. Deeper ponds and lakes may thermally stratify in the summer months and result in deeper waters becoming depleted of oxygen. Lake stratification is a result of the peculiar property of water density changes with temperature. The density of all liquid changes with changes in temperature, however, water behaves in a special way. As most liquids are cooled the density, or relative weight, of the liquid increases due to the compaction of the molecules in the liquid, and conversely, as liquids are heated the density decreases. Water, because of its unique characteristics, is at its maximum density at 4 degrees Centigrade, or approximately 39.2 degrees Fahrenheit. When water is either heated above this temperature or cooled below this temperature its density decreases. This is why ice floats, or forms on the surface of lakes and ponds. A normal cycle of stratification in lakes in our region of the country, in early spring after ice out is as follows: the lake water will all be nearly the same temperature shortly after ice out and wind action on the lake surface will induce circulation of the entire volume of water. As spring advances and the increased sunlight energy warms the surface waters, these become lighter and tend to separate from the deeper waters and essentially float on top of the cooler water. This continues until there is a very stable "layering" or stratification of water in the lake. There are three distinct layers of water in stratified lakes, as described by limnologists:

1. Epilimnion (upper warm layer) - this is, generally speaking, confined to the top 10 ft. to 15 ft. of the lake volume. This is a warm region, mixed thoroughly by wind to a more or less uniform temperature. It is also the zone of most photosynthetic production and is usually high in dissolved oxygen.
2. Thermocline or Metalimnion (middle layer of rapidly changing temperature) this layer is the area in the lake where temperature decreases rapidly, usually about 1 degree centigrade or more per meter (or approximately 3 ft.). Oxygen depletion generally begins in this layer.
3. Hypolimnion (deep, cold layer) - this layer is relatively unaffected by wind mixing or motor boat activity, and is often devoid of oxygen. Oxygen is depleted by the decomposition of dead organic matter falling from the upper waters as well as external sources such as leaves and grass clippings that sink to the bottom of the lake.

Once this stratification is established (usually by early to mid-June, in our area) it is very stable and stays intact until the fall turnover, which is caused by decreasing surface water temperatures (causing increased density), and the mixing of the lake waters by the wind. The lake then circulates completely for a period of time, usually until ice cover forms, sealing off the surface of the lake from the atmosphere. A reverse stratification then sets in where the water just under the ice is just above 32 degrees Fahrenheit with increasing temperature with depth to a temperature of 39.2 degrees Fahrenheit. Decomposition continues in the bottom throughout the winter, resulting in oxygen depletion in the bottom waters. This progresses towards the surface throughout ice cover and can cause an oxygen depletion fish kill under the ice during severe winters. After the ice melts, the lake begins to circulate again, and the cycle has completed itself. This phenomenon has a profound affect on the biological and chemical components of the lake ecosystem. Alkalinity is the ability of water to buffer against pH changes upon the addition of an acid or base. The alkalinity of a lake or pond is generally determined by the characteristics of the watershed or local geology. As a general guideline a well-buffered system has an alkalinity of 50 parts per million (ppm) or greater. Well buffered systems have potential for moderate to high productivity. Alkalinity is important in determining algacide dosages, particularly copper sulfate. The maximum safe dosage for fish of copper sulfate if total alkalinity is less than 50 ppm is 0.25 ppm or .68 pounds / acre-foot, 1.00 ppm or

2.7 pounds / acre-foot for a total alkalinity range of 50 to 200 ppm, and 1.5 ppm or 4.0 pounds / acre- foot for a total alkalinity greater than 200 ppm.

Hardness is a measure of the calcium and magnesium (and some other ions) concentrations in water. The concept of hardness comes from the field of domestic water supply. It is a measure of soap requirements for adequate lather formation and is an indicator of the rate of scale formation in hot water heaters. Hardness and alkalinity are sometimes used interchangeably; however, these parameters sometimes have very different values. Waters containing a hardness measure of greater than 75 ppm may be considered hard and are often clearer and weedier than soft waters (Walker et al., 1985).

Nitrogen can exist in several forms within a body of water, including: ammonia, nitrite, nitrate, and organic nitrogen (amino acids and proteins). Ammonia results from the biological decomposition of organic matter by bacteria. During the process of nitrification, nitrate (which is directly available for plant uptake) is formed from the complete biological oxidation of ammonia in which nitrite is an intermediate product. Nitrate is a major plant nutrient. The most important forms of nitrogen for the growth of algae include ammonia and nitrate. Total nitrogen levels above 1.9 ppm are generally indicative of nutrient enrichment or eutrophic conditions (Wetzel, 1983). Inorganic nitrogen (nitrite, nitrate, ammonia, and ammonium) levels greater than 0.30 ppm are indicative of eutrophic lakes and ponds (Sawyer, 1948). Nitrogen in surface waters cannot be controlled by any economical method. Blue-green algae can fix nitrogen directly from the atmosphere unlike other forms of plants.

Phosphorus is a major plant nutrient and is most often the limiting factor for algae and macrophyte (vascular plants) growth within an aquatic system. Total phosphorus levels in excess of 0.03 ppm indicate eutrophic conditions (Vollenwieder, 1968). Waters with excessive plant growth high nutrients and degraded water quality are typical of eutrophic lakes and ponds. Ortho-phosphorus is a measure of the dissolved inorganic phosphorus available for immediate plant uptake. Concentrations of ortho-phosphate above 0.045 ppm may be considered critical concentrations above which nuisance algae blooms could be expected (Sawyer, 1948). The remainder of the total phosphorus is most likely bound onto particulate material and although not immediately available for uptake, could become available through biochemical degradation. Dissolved phosphorus is absorbed from the water column primarily by phytoplankton. Phosphorus supply to aquatic macrophytes is primarily from the sediment rather than from the water column.

Phosphorus is released from sediment under anaerobic conditions but is precipitated to the sediment under aerobic conditions. Phosphorus can be precipitated from the water column by use of alum (aluminum sulfate). Sediment phosphorus can be inactivated and made unavailable to macrophytes by heavy applications of alum to the sediment surface.

EQUIPMENT AND METHODS

Water quality analysis equipment used in this survey included a YSI ProODO oxygen-temperature meter with 60 ft. remote sensing probe, a Hach field test kit, and a Wildco Alpha Water bottle sampler. The following water quality parameters were measured and recorded: dissolved oxygen, temperature, pH, total hardness, total alkalinity, nitrate-nitrogen, and orthophosphate. The parameters selected are the major water quality factors influencing the lakes productivity and fish health. Water quality testing to determine nutrient levels was completed in the lab using a Hach DR/2010 photospectrometer.

Fish sampling was done with the use of an electrofishing boat. Electrofishing is simply the use of electricity to capture fish for the evaluation of population status. Various types of equipment are in use today, however, most fisheries biologists agree that pulsed DC current is more efficient and less harmful to the fish collected than AC current.

Electrofishing with an experienced crew using proven equipment is probably the least selective method of sampling warm water fish species in the temperate waters of our area. Evaluation of electrofishing efficiencies have shown that night electrofishing is a reliable method for sampling populations of largemouth bass, bluegill, and redear sunfish, and generally detects the presence of green sunfish and other scaled fish species. The method is less efficient for sampling populations of channel catfish, bullheads, and crappie (Reynolds and Simpson, 1976). The largest bias in electrofishing is generally that of collecting more large fish of a given species than smaller individuals. This is due to the differential effect of the electric current on fish of different sizes, interference with collection from dense weed beds, if present, as well as the potential bias of the person dipping stunned fish (Nielsen & Johnson, 1983). Many years of experience by our personnel has reduced this bias to an acceptable level.

Electrofishing equipment used in this survey consisted of a 16 foot workboat equipped with a Midwest Lake Electrofishing Systems Infinity Box powered by a 6500 watt portable generator and a boom mounted electrosphere designed by Coffelt Manufacturing. The electrosphere allows the use of higher voltages at lower amperage. This has proven

to improve the efficiency of the electrofishing technique with lower damage to captured fish. The electrofisher was operated in the pulsed DC mode using an output level of 350 to 375 volts. The increased effectiveness of this electrofishing system makes fish of all species, including channel catfish, more vulnerable to capture. This results in a better sampling of all fish species with a higher capture rate of the more vulnerable species (bass, bluegill, redear, and green sunfish) in the samples taken. All fish collected were placed in water filled containers aboard the sampling boat for processing. A sub-sample of up to five specimens from each species was taken in each one-half inch group. The individual fish in these sub samples were weighed to the nearest hundredth pound for average weight determinations. Additional specimens were recorded by length group.

Field data was summarized and is presented in table and graph form. Condition factors and relative weight calculations (standard measurements of the relative plumpness) were calculated for important species using standard formulas (Anderson and Gutreuter, Carlander 1977, Hillman 1982, Wege and Anderson 1978). Relative weight is a comparison of fish weights at certain sizes to standard calculated weights at those sizes. Relative weights of 100 or greater are considered good. An important procedure used in our evaluation of the bass – bluegill populations, and other species to a lesser extent, is the Proportional Stock Density Index. This population index was developed by intensive research into dynamics of fish population structure, primarily in largemouth bass - bluegill dominated populations (Anderson 1976), and subsequent field testing by numerous fisheries research and management biologists in mid-western states. Bluegill samples are divided into three major groups: those less than 3.0 inches in length, those 3.0 inches and larger, and those 6.0 inches and larger. The group 3.0 inches and larger are called the "stock". The 6.0-inch and larger individuals are considered to be "quality" or harvestable size. Bluegill PSD is the percentage of bluegill "stock" that is in the harvestable size. Largemouth bass samples are separated into "stock size" (8.0 inches plus) and quality size (12.0 inches plus), for PSD calculations. Largemouth bass PSD is the percentage of bass "stock" that are of a "quality" or harvestable size. This study, and subsequent studies and application of the techniques developed in those studies, have shown that fish populations that are producing, or are capable of producing, a sustained annual harvest of "quality" largemouth bass and bluegill have certain characteristics. These include the following:

1. Reasonably high numbers of bluegill smaller than 2.5 inches (young-of-the-year)

2. Proportional Stock Density index of 20 - 40 for bluegill.
3. Bluegill growth which results in a length of 6.0 inches by age III or IV, with low numbers of age V or older fish.
4. Proportional Stock Density index of 40 - 60 for largemouth bass.
5. A minimum of 20 adult bass per acre.
6. A maximum of 50% annual mortality (harvest) of bass in age II - V.
7. Growth rate that results in 8 inch bass reaching quality size (12 inch plus) in approximately 1 year.
8. No missing year classes in ages 0 - V.
9. A maximum of 10% of the lake bottom covered by dense weed beds.

These parameters, and other factors, are used in the evaluation and development of recommendations from Aquatic Control surveys.

LITERATURE CITED AND REFERENCE LIST

- Anderson, R. 1973. Applications of theory and research to management of warmwater fish populations. Trans. Am. Fish. Soc. 102(1)164-171.
- Anderson, R. 1976. Management of small warmwater impoundments. Fisheries 1(6): 5-7, 26-28.
- Anderson, R., and S.J. Gutreuter. 1983. Length, weight, and associated structural indices Pages 283-300 in L. A. Nielsen and D. L. Johnson, editors. Fisheries Techniques. American Fisheries Society, Bethesda, Maryland.
- Arnold, D.E. 1971. Ingestion, assimilation, survival, and reproduction by *Daphnia pulex* fed seven species of blue-green algae. Limnology and Oceanography. 16: 906-920.
- Bennett, C. W. 1971. Management of lakes and ponds. Van Nostrand Reinhold. G. New York 375 pp.
- Boyd, C.E. 1990. Water quality in ponds for Aquaculture. Auburn Univ. Ag. Exp. Sta. Auburn, Al. 252 pp.
- Calhoun, A. (editor) 1966. Inland Fisheries Management. State of California. Dept. of Fish & Game, 546 pp.
- Carlander, K. D. 1969 & 1977. Handbook of freshwater fishery biology. Vols. 1 & 2. Iowa State University Press, Ames, Iowa, Vol 1. 752 pp, Vol 2, 409 pp.

- Cole, Gerald, A. 1983. Textbook of Limnology. 3 ed. Dept. of Zoology, Arizona State Univ. Tempe, AZ. The C.V. Mosby Co. St. Louis.
- D'Itri, F. (editor) 1985. Artificial reefs Marine and Freshwater applications, Lewis Publishers, Inc. Chelsea, MI 589 pp.
- Funk, J. L. (editor) 1974. Symposium on overharvest and management of largemouth bass in small impoundments. North Central Div. Am.Fish. Soc. Sp. Publ. No. 3 116 pp.
- Hayes, J. W., and T. E. Wissing. 1996. Effects of stem density of artificial vegetation on abundance and growth of age-0 bluegills and predation by largemouth bass. Transactions of the American Fisheries Society 125:422-433
- Hillman, W.P. 1982. Structure and dynamics of unique bluegill populations. Master's Thesis. University of Missouri, Columbia.
- Indiana Dept of Nat. Res. 1966, 1985, 1988, Guidelines for the evaluation of sport fish populations in Indiana. Unpublished data.
- Johnson, D.L. & Stein, R.A. 1979. (editors) Response of fish to habitat structure in standing water. North Cen. Am. Fish Soc. Sp. Publ. No. 6. 77pp.
- Kornman, L.E. 1990. Evaluation of a 15-inch minimum size limit on Black Bass at Grayson Lake, Bull. #90. State of KY Dept. of Fish & Wildlife Res. 71pp.
- Kwak, T. J., M. G. Henry. 1995. Largemouth bass mortality and related causal factors During live release fishing tournaments on a large Minnesota lake. North American Journal of Fisheries Management 15: 621-630.
- Lawrence, J.M. 1958. Estimated size of various forage fishes largemouth bass can swallow. Proc. of 11th Annual Conf. S.E. Assoc. Fish & Game Comm. 220-225.
- Lyons, John. Fish of Wisconsin Identification Database. Picture of Gizzard Shad. 30 June 2004. University of Wisconsin Center for Limnology, Wisconsin Sea Grant, Wisconsin Dept. of Natural Resources. 12 Nov. 2015.
< <http://www.seagrant.wisc.edu/home/Default.aspx?tabid=605&FishID=56>>
- Lyons, John. Fish of Wisconsin Identification Database. Picture of Black Crappie. 30 June 2004. University of Wisconsin Center for Limnology, Wisconsin Sea Grant, Wisconsin Dept. of Natural Resources. 12 Nov. 2015.
< <http://www.seagrant.wisc.edu/home/Default.aspx?tabid=605&FishID=11>>
- Lyons, John. Fish of Wisconsin Identification Database. Picture of Brown Bullhead. 30 June 2004. University of Wisconsin Center for Limnology, Wisconsin Sea Grant, Wisconsin Dept. of Natural Resources. 12 Nov. 2015.

< <http://www.seagrant.wisc.edu/home/Default.aspx?tabid=605&FishID=29>>

Lyons, John. Fish of Wisconsin Identification Database. Picture of Bluntnose Minnow.
30 June 2004. University of Wisconsin Center for Limnology, Wisconsin Sea Grant, Wisconsin Dept. of Natural Resources. 12 Nov. 2015.

< <http://www.seagrant.wisc.edu/home/Default.aspx?tabid=605&FishID=23>>

Lyons, John. Fish of Wisconsin Identification Database. Picture of Common Carp.
30 June 2004. University of Wisconsin Center for Limnology, Wisconsin Sea Grant, Wisconsin Dept. of Natural Resources. 12 Nov. 2015.

< <http://www.seagrant.wisc.edu/home/Default.aspx?tabid=605&FishID=33>>

Lyons, John. Fish of Wisconsin Identification Database. Picture of White Sucker.
30 June 2004. University of Wisconsin Center for Limnology, Wisconsin Sea Grant, Wisconsin Dept. of Natural Resources. 12 Nov. 2015.

< <http://www.seagrant.wisc.edu/home/Default.aspx?tabid=605&FishID=163>>

McComas, S. 1993. Lake Smarts The First Lake Maintenance Handbook. Terrene Institute, Washington, D.C. 215pp.

Mittelbach, G. G. 1981. Foraging efficiency and body size: a study of optimal diet and Habitat use by bluegills. Ecology 65:1370-1386

National Academy of Sci. 1969. Eutrophication, causes, consequences, correctives. Washington D.C. 658pp.

Nielsen, L.A. and Johnson, D.L. (editors) 1983. Fisheries Techniques. Am. Fish. Soc. Southern Printing Co., Inc. Blacksburg, VA. 468 pp.

Novinger, G.D. & Dillard, J. 1978. New approaches to the management of small impoundments. North Cen. Div. Am. Fish. Soc. Sp. Publ. No. 5. 132 pp.

Pereira, D.L., S.A. Pothaven, and B. Vondracek. 1999. Effects of Vegetation Removal on Bluegill and Largemouth Bass in Two Minnesota Lakes. North American Journal of Fisheries Management 19: 748-756.

Pflieger, W. L. 1975. The Fishes of Missouri. Missouri Department of Conservation. 343pp.

Prather, K.W. 1990. Evaluation of a 12-16 Inch Slot limit on largemouth bass at Elmer Davis Lake. State of KY. Dept. of Fish & Wildlife Res. Bull. #89. 18pp

Reynolds & Simpson. 1976. Evaluation of fish sampling methods and rotenone census. pages in: Novinger & Dillard. 1978. New approaches to the management of small impoundments. N.C. Div. Am. Fish. Soc. Sp. Publ. No. 5 132 pp.

- Ruttner, Franz. 1953. Fundamentals of limnology. 3rd edition. Univ. of Toronto Press. Toronto. 261pp.
- Sawyer, C. N. 1948. Fertilization of Lakes by Agricultural and Urban Drainage. Journal of the New England Water Works Association, 61 109-127.
- Savino, J.F., and R.A. Stein. 1982 Predator-prey interactions between largemouth bass and bluegills as influenced by simulated, submerged vegetation. Transactions of the American Fisheries Society 111: 255-256 Sport Fishing Inst. 1975. Black Bass Biology & Management. Washington. D.C. 534pp.
- Strange, R. J., C. R. Berry, and C. B. Schreck. 1975. Aquatic Plant control and reservoir fisheries. Pages 513-521 in R. H. Stroud, editor. Predator-prey systems in fisheries management. Sport Fishing Institute, Washington D.C.
- Taras, M. J., A. E. Greenberg, R. D. Hoak, and M. C. Rand eds. 1971. Standard Methods for the Examination of Water and Wastewater. American Public Health Association, Washington D.C. 874pp.
- U S E.P.A. 1976. Quality Criteria for Water. U.S. Govt. Printing Office. 256 pp.
- Vollenweider, R. A. 1968. Scientific Fundamentals of the Eutrophication of Lakes and Flowing Waters, with Particular Reference to Nitrogen and Phosphorous as Factors in Eutrophication. OECD Report No. DAS/CSI/68.27, Paris.
- Wege & Anderson. 1978. Relative Weight(Wr): A new Index of condition for largemouth bass. pages in: Novinger & Dillard. 1978. New approaches to the management of small impoundments. N.C. Div. Am. Fish Soc. Sp. Publ. No. 5. 132pp.
- Werner, E.E., and D.J. Hall. 1988. Ontogenetic niche shifts in bluegill: the foraging rate predation risk trade-off. Ecology 69:1352-1366
- Wiley, M. J. W. Gorden, S. W. Waite, and T. Powless. 1984. The relationship between aquatic macrophytes and sport fish production in Illinois ponds: a simple model. North American Journal of Fisheries Management 4:111-119.

Appendix B

Fish Collection Table

Size Group (IN)	NUMBER	PERCENTAGE	AVERAGE WEIGHT (lbs.)	TOTAL WEIGHT (lbs.)	CONDITION FACTOR	WS	RELATIVE WEIGHT
-----------------------	--------	------------	-----------------------------	---------------------------	---------------------	----	--------------------

BLUEGILL

<3.0	90	8.08%	0.01	0.90	-	-	-
3.0	24	2.15%	0.02	0.43	6.67	0.02	111
3.5	5	0.45%	0.03	0.17	7.93	0.03	125
4.0	11	0.99%	0.04	0.40	5.63	0.04	85
4.5	55	4.94%	0.06	3.19	6.36	0.06	93
5.0	71	6.37%	0.07	4.62	5.20	0.09	73
5.5	75	6.73%	0.10	7.65	6.13	0.12	84
6.0	234	21.01%	0.13	29.95	5.93	0.16	79
6.5	253	22.71%	0.16	39.47	5.68	0.21	74
7.0	270	24.24%	0.20	54.00	5.83	0.27	74
7.5	26	2.33%	0.25	6.45	5.88	0.34	73
TOTAL		1114		147.22			

GIZZARD SHAD

4.0	3	0.73%	0.04	0.11
4.5	8	1.96%	0.03	0.26
5.0	52	12.71%	0.05	2.60
5.5	22	5.38%	0.06	1.23
6.0	3	0.73%	0.06	0.17
9.5	9	2.20%	0.33	2.93
10.0	12	2.93%	0.34	4.10
10.5	26	6.36%	0.43	11.08
11.0	61	14.91%	0.47	28.79
11.5	64	15.65%	0.54	34.82
12.0	62	15.16%	0.65	40.30
12.5	37	9.05%	0.70	25.90
13.0	28	6.85%	0.72	20.10
13.5	12	2.93%	0.89	10.63
14.0	6	1.47%	0.95	5.71
15.0	3	0.73%	1.48	4.44
15.5	1	0.24%	1.33	1.33
TOTAL		409		194.50

BLACK CRAPPIE

4.0	4	2.47%	0.03	0.13
4.5	1	0.62%	0.04	0.04
6.0	1	0.62%	0.10	0.10
6.5	4	2.47%	0.12	0.47
7.0	28	17.28%	0.15	4.20
7.5	82	50.62%	0.19	15.25
8.0	39	24.07%	0.20	7.96
8.5	2	1.23%	0.24	0.48
9.5	1	0.62%	0.34	0.34
TOTAL		162		28.97

**CHANNEL
CATFISH**

4.0	1	0.78%	0.04	0.04
6.0	1	0.78%	0.09	0.09
7.0	1	0.78%	0.14	0.14
8.0	6	4.69%	0.26	1.58
8.5	17	13.28%	0.32	5.44
9.0	52	40.63%	0.30	15.50
9.5	22	17.19%	0.39	8.62
10.0	11	8.59%	0.45	4.95
10.5	2	1.56%	0.54	1.08
11.5	1	0.78%	0.41	0.41
13.0	1	0.78%	0.60	0.60
13.5	2	1.56%	0.81	1.61
14.0	3	2.34%	0.85	2.55
14.5	1	0.78%	0.82	0.82
15.5	1	0.78%	1.40	1.40
16.5	1	0.78%	1.50	1.50
17.0	1	0.78%	1.70	1.70
18.5	1	0.78%	2.50	2.50
19.0	1	0.78%	2.14	2.14
22.0	1	0.78%	3.75	3.75
23.0	1	0.78%	5.15	5.15
TOTAL		128	61.57	

**LARGEMOUTH
BASS**

3.0	1	1.00%	0.02	0.02	7.41	0.01	-
3.5	2	2.00%	0.03	0.06	7.00	0.02	-
4.0	4	4.00%	0.03	0.11	4.30	0.03	-
4.5	2	2.00%	0.04	0.08	4.39	0.04	-
5.0	2	2.00%	0.06	0.11	4.40	0.06	-
5.5	1	1.00%	0.06	0.06	3.61	0.07	-
6.5	1	1.00%	0.13	0.13	4.73	0.13	-
7.0	1	1.00%	0.15	0.15	4.37	0.16	-
7.5	3	3.00%	0.16	0.49	3.86	0.20	-
8.0	5	5.00%	0.18	0.92	3.59	0.25	75
8.5	3	3.00%	0.26	0.78	4.23	0.30	87
9.0	8	8.00%	0.32	2.59	4.44	0.36	90
9.5	11	11.00%	0.40	4.40	4.67	0.43	94
10.0	7	7.00%	0.46	3.23	4.62	0.50	92
10.5	3	3.00%	0.56	1.68	4.84	0.59	95
11.0	3	3.00%	0.66	1.97	4.93	0.68	96
11.5	1	1.00%	0.69	0.69	4.54	0.78	88
12.0	2	2.00%	0.86	1.72	4.98	0.90	96
12.5	1	1.00%	0.93	0.93	4.76	1.02	91
13.0	2	2.00%	1.13	2.25	5.12	1.16	97
13.5	1	1.00%	1.10	1.10	4.47	1.31	84
14.0	5	5.00%	1.39	6.95	5.07	1.47	95
14.5	1	1.00%	1.80	1.80	5.90	1.64	109
15.0	3	3.00%	1.83	5.48	5.41	1.83	100
15.5	3	3.00%	1.74	5.22	4.67	2.03	86
16.0	5	5.00%	2.24	11.22	5.48	2.25	100
16.5	5	5.00%	2.30	11.52	5.13	2.48	93

17.0	6	6.00%	2.57	15.41	5.23	2.73	94
18.0	3	3.00%	2.14	6.43	3.68	3.28	65
18.5	1	1.00%	3.25	3.25	5.13	3.58	91
19.0	4	4.00%	3.91	15.64	5.70	3.89	100
TOTAL				106.39			

**BROWN
BULLHEAD**

7.5	1	2.00%	0.21	0.21
8.0	1	2.00%	0.26	0.26
8.5	3	6.00%	0.29	0.87
9.0	24	48.00%	0.30	7.15
9.5	12	24.00%	0.40	4.75
10.0	4	8.00%	0.45	1.79
10.5	4	8.00%	0.50	1.98
11.0	1	2.00%	0.49	0.49
TOTAL				17.50

GREEN SUNFISH

<3.0	4	21.05%	0.01	0.04
4.0	2	10.53%	0.04	0.07
4.5	5	26.32%	0.06	0.28
5.0	4	21.05%	0.10	0.41
5.5	1	5.26%	0.10	0.10
6.0	3	15.79%	0.13	0.39
TOTAL				1.29

HYBRID STRIPED BASS

10.5	1	25.00%	0.48	0.48
11.0	1	25.00%	0.49	0.49
13.5	1	25.00%	0.77	0.77
15.0	1	25.00%	1.50	1.50
TOTAL				3.24

BLUNTNOST MINNOW

<3.0	4	100.00%	0.01	0.04
TOTAL				0.04

REDEAR SUNFISH

9.0	1	50.00%	0.65	0.65
11.0	1	50.00%	1.24	1.24
TOTAL				1.89

COMMON CARP

29.0	1	50.00%	0.26	13.50
33.5	1	50.00%	0.73	14.50
TOTAL				28.00

WHITE SUCKER

15.0	1	50.00%	1.45	1.45
17.5	1	50.00%	2.31	2.31
TOTAL				3.76

